

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Biextensions and 1-motives

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/138511> since 2018-03-23T15:33:34Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

BIEXTENSIONS OF 1-MOTIVES BY 1-MOTIVES

CRISTIANA BERTOLIN

ABSTRACT. Let S be a scheme. In this paper, we define the notion of biextensions of 1-motives by 1-motives. Moreover, if $\mathcal{M}(S)$ denotes the Tannakian category generated by 1-motives over S (in a geometrical sense), we define geometrically the morphisms of $\mathcal{M}(S)$ from the tensor product of two 1-motives $M_1 \otimes M_2$ to another 1-motive M_3 , to be the isomorphism classes of biextensions of (M_1, M_2) by M_3 :

$$\mathrm{Hom}_{\mathcal{M}(S)}(M_1 \otimes M_2, M_3) = \mathrm{Biext}^1(M_1, M_2; M_3)$$

Generalizing this definition we obtain, modulo isogeny, the geometrical notion of morphism of $\mathcal{M}(S)$ from a finite tensor product of 1-motives to another 1-motive.

INTRODUCTION

Let S be a scheme.

A 1-motive over S consists of a S -group scheme X which is locally for the étale topology a constant group scheme defined by a finitely generated free \mathbb{Z} -module, a semi-abelian S -scheme G , and a morphism $u : X \rightarrow G$ of S -group schemes.

Let $\mathcal{M}(S)$ be what should be the Tannakian category generated by 1-motives over S in a geometrical sense. We know very little about this category $\mathcal{M}(S)$: in particular, we are not able to describe geometrically the object of $\mathcal{M}(S)$ defined as the tensor product of two 1-motives! Only if $S = \mathrm{Spec}(k)$, with k a field of characteristic 0 embeddable in \mathbb{C} , we know something about $\mathcal{M}(k)$: in fact, identifying 1-motives with their mixed realizations, we can identify $\mathcal{M}(k)$ with the Tannakian sub-category of an “appropriate” Tannakian category of mixed realizations generated by the mixed realizations of 1-motives.

The aim of this paper is to use biextensions in order to define some morphisms in the category $\mathcal{M}(S)$: *Geometrically the morphisms of $\mathcal{M}(S)$ from the tensor product of two 1-motives $M_1 \otimes M_2$ to another 1-motive M_3 are the isomorphism classes of biextensions of (M_1, M_2) by M_3 :*

$$(0.0.1) \quad \mathrm{Hom}_{\mathcal{M}(S)}(M_1 \otimes M_2, M_3) = \mathrm{Biext}^1(M_1, M_2; M_3)$$

The idea of defining morphisms through biextensions goes back to Alexander Grothendieck. In fact, in [6] Exposé VIII he defines pairings from biextensions: if P, Q, G are three abelian groups of a topos \mathbf{T} , to each isomorphism class of biextensions of (P, Q) by G , he associates a pairing $({}_l P)_{n \geq 0} \otimes ({}_l Q)_{n \geq 0} \rightarrow ({}_l G)_{n \geq 0}$ where $({}_l P)_{n \geq 0}$ (resp. $({}_l Q)_{n \geq 0}$, $({}_l G)_{n \geq 0}$) is the projective system constructed from the kernels ${}_l P$ (resp. ${}_l Q$, ${}_l G$) of the multiplication by l^n for each $n \geq 0$.

1991 *Mathematics Subject Classification.* 18A20;14A15.

Key words and phrases. biextensions, 1-motives, tensor products, morphisms.

Generalizing Grothendieck's work, in "Theorie de Hodge III" Pierre Deligne defines the notion of biextension of two complexes of abelian groups concentrated in degree 0 and -1 (over any topos \mathbf{T}) by an abelian group. Applying this definition to 1-motives and to \mathbb{G}_m , to each isomorphism class of such biextensions he associates a pairing $T_*(M_1) \otimes T_*(M_2) \longrightarrow T_*(\mathbb{G}_m)$ in the Hodge, De Rham, ℓ -adic realizations (resp. $*$ = H, $*$ = dR, $*$ = ℓ).

Our definition of biextension of 1-motives by 1-motive generalizes the one of Deligne. The main idea of our definition is the following: Recall that a 1-motive M over S can be described also as a 7-uplet $(X, Y^\vee, A, A^*, v, v^*, \psi)$ where

- X and Y^\vee are two S -group schemes which are locally for the étale topology constant group schemes defined by finitely generated free \mathbb{Z} -modules;
- A and A^* are two abelian S -schemes dual to each other;
- $v : X \longrightarrow A$ and $v^* : Y^\vee \longrightarrow A^*$ are two morphisms of S -group schemes; and
- ψ is a trivialization of the pull-back $(v, v^*)^* \mathcal{P}_A$ via (v, v^*) of the Poincaré biextension \mathcal{P}_A of (A, A^*) .

In other words, we can reconstruct the 1-motive M from the Poincaré biextension \mathcal{P}_A of (A, A^*) and some trivializations of some pull-backs of \mathcal{P}_A . The 4-uplet (X, Y^\vee, A, A^*) corresponds to the pure part of the 1-motive, i.e. it defines the pure motives underlying M , and the 3-uplet v, v^*, ψ represents the "mixity" of M . Therefore the Poincaré biextension \mathcal{P}_A is related to the pure part (X, Y^\vee, A, A^*) of the 1-motive and the trivializations are related to the mixed part v, v^*, ψ of M .

Now let $M_i = [X_i \xrightarrow{u_i} G_i]$ (for $i = 1, 2, 3$) be a 1-motive over S . A biextension of (M_1, M_2) by M_3 is a biextension \mathcal{B} of (G_1, G_2) by G_3 , some trivializations of some pull-backs of \mathcal{B} , and a morphism $X_1 \otimes X_2 \longrightarrow X_3$ compatible with the above trivializations. Since by the main Theorem of [2], to have a biextension of (G_1, G_2) by G_3 is the same thing has to have a biextension of (A_1, A_2) by $Y_3(1)$, we can simplify our definition: a biextension $B = (B, \Psi_1, \Psi_2, \Psi, \lambda)$ of (M_1, M_2) by M_3 consists of

- (1) a biextension of B of (A_1, A_2) by $Y_3(1)$;
- (2) a trivialization Ψ_1 (resp. Ψ_2) of the biextension $(v_1, id_{A_2})^* B$ (resp. $(id_{A_1}, v_2)^* B$) of (X_1, A_2) by $Y_3(1)$ (resp. of (A_1, X_2) by $Y_3(1)$) obtained as pull-back of the biextension B via (v_1, id_{A_2}) (resp. via (id_{A_1}, v_2));
- (3) a trivialization Ψ of the biextension $(v_1, v_2)^* B$ of (X_1, X_2) by $Y_3(1)$ obtained as pull-back of the biextension B via (v_1, v_2) , which coincides with the trivializations induced by Ψ_1 and Ψ_2 over $X_1 \times X_2$;
- (4) a morphism $\lambda : X_1 \otimes X_2 \longrightarrow X_3$ compatible with the trivialization Ψ and the morphism $u_3 : X_3 \longrightarrow G_3$.

The biextension B and the morphism $\lambda : X_1 \otimes X_2 \longrightarrow X_3$ take care of the pure parts of the 1-motives M_1, M_2 and M_3 , i.e. of X_i, Y_i, A_i, A_i^* for $i = 1, 2, 3$. On the other hand the trivializations Ψ_1, Ψ_2, Ψ , and the compatibility of λ with such trivializations and with u_3 , take care of the mixed parts of M_1, M_2 and M_3 , i.e. of v_i, v_i^* and ψ_i for $i = 1, 2, 3$.

In terms of morphisms, the biextension $B = (B, \Psi_1, \Psi_2, \Psi, \lambda)$ defines a morphism $M_1 \otimes M_2 \longrightarrow M_3$ where

- the biextension B of (A_1, A_2) by $Y_3(1)$ defines the component $A_1 \otimes A_2 \longrightarrow Y_3(1)$;

- the morphism λ defines the component $X_1 \otimes X_2 \longrightarrow X_3$ of weight 0;
- the trivializations Ψ_1, Ψ_2, Ψ and the compatibility of λ with such trivializations and with u_3 take care of the other components of $M_1 \otimes M_2 \longrightarrow M_3$, and of the compatibility of the mixed part of $M_1 \otimes M_2$ and the mixed part of M_3 through this morphism of $M_1 \otimes M_2 \longrightarrow M_3$.

At the end of Section 1, we give several examples of biextensions of 1-motives by 1-motives. At the end of Section 2, we show how to construct explicitly the morphisms corresponding to the biextensions stated as example at the end of Section 1.

Then we verify that the property of *respecting weights* which is satisfied by the morphisms of $\mathcal{M}(S)$, is also verified by biextensions. For example, in $\mathcal{M}(S)$ there are no morphisms from $X_1 \otimes X_2$ to G_3 or from $G_1 \otimes G_2$ to X_3 (Lemma 2.1.1), and therefore we must have that all the biextensions of $([X_1 \rightarrow 0], [X_2 \rightarrow 0])$ by $[0 \rightarrow G_3]$ and all the biextensions $([0 \rightarrow G_1], [0 \rightarrow G_2])$ by $[X_3 \rightarrow 0]$ are trivial. To have a morphism from $G_1 \otimes G_2$ to G_3 is the same thing as to have a morphism from $A_1 \otimes A_2$ to $Y_3(1)$ (Corollary 2.1.3) and therefore we must have that to have a biextension of (G_1, G_2) by G_3 is the same thing as to have a biextension of (A_1, A_2) by $Y_3(1)$ (cf. Theorem [2]).

We can extend definition (0.0.1) to a finite tensor product of 1-motives in the following way: again because of weights, a morphism from a finite tensor product $\otimes_1^l M_j$ of 1-motives to a 1-motive M involves only the quotient $\otimes_1^l M_j / W_{-3}(\otimes_1^l M_j)$ of the mixed motive $\otimes_1^l M_j$. However the motive $\otimes_1^l M_j / W_{-3}(\otimes_1^l M_j)$ is isogeneous to a finite sum of copies of $M_{\iota_1} \otimes M_{\iota_2}$ for $\iota_1, \iota_2 \in \{1, \dots, l\}$ (Lemma 2.2.6), and therefore, modulo isogeny, a morphism from a finite tensor product of 1-motives to a 1-motive is a sum of isomorphism classes of biextensions of 1-motives by 1-motives (Theorem 2.2.7).

A special case of definition (0.0.1) was already used in the computation of the unipotent radical of the Lie algebra of the motivic Galois group of a 1-motive defined over a field k of characteristic 0 (cf. [1]). In fact in [1] (1.3.1), using Deligne's definition of biextension of 1-motives by \mathbb{G}_m , we define a morphism from the tensor product $M_1 \otimes M_2$ of two 1-motives to a torus as an isomorphism class of biextensions of (M_1, M_2) by this torus.

ACKNOWLEDGMENT

I want to thank the referee of my article [1] who suggested me to generalize it: the study of biextensions of 1-motives by 1-motives started with the attempt of generalizing the definition [1] (1.3.1).

In this paper S is a scheme.

1. BIEXTENSIONS OF 1-MOTIVES BY 1-MOTIVES

In [3] (10.1.10), Deligne defines a **1-motive** M over S as

- (1) a S -group scheme X which is locally for the étale topology a constant group scheme defined by a finitely generated free \mathbb{Z} -module,
- (2) a semi-abelian S -scheme G , i.e. an extension of an abelian S -scheme A by a S -torus $Y(1)$, with cocharacter group Y ,
- (3) a morphism $u : X \longrightarrow G$ of S -group schemes.

The 1-motive M can be view also as a complex $[X \xrightarrow{u} G]$ of commutative S -group schemes concentrated in degree 0 and -1. An **isogeny between two 1-motives** $M_1 = [X_1 \xrightarrow{u_1} G_1]$ and $M_2 = [X_2 \xrightarrow{u_2} G_2]$ is a morphism of complexes (f_X, f_G) such that $f_X : X_1 \rightarrow X_2$ is injective with finite cokernel, and $f_G : G_1 \rightarrow G_2$ is surjective with finite kernel.

1-motives are mixed motives of niveau ≤ 1 : the weight filtration W_* on $M = [X \xrightarrow{u} G]$ is

$$\begin{aligned} W_i(M) &= M \text{ for each } i \geq 0, \\ W_{-1}(M) &= [0 \rightarrow G], \\ W_{-2}(M) &= [0 \rightarrow Y(1)], \\ W_j(M) &= 0 \text{ for each } j \leq -3. \end{aligned}$$

In particular, we have $\text{Gr}_0^W(M) = [X \rightarrow 0]$, $\text{Gr}_{-1}^W(M) = [0 \rightarrow A]$ and $\text{Gr}_{-2}^W(M) = [0 \rightarrow Y(1)]$.

1.1. The category of biextensions of 1-motives by 1-motives. Let $M_i = [X_i \xrightarrow{u_i} G_i]$ (for $i = 1, 2, 3$) be a 1-motive over S . The following definition of biextension of (M_1, M_2) by M_3 , is a generalization of Deligne's definition [3] (10.2):

Definition 1.1.1. A biextension $\mathcal{B} = (\mathcal{B}, \Psi_1, \Psi_2, \Psi, \lambda)$ of (M_1, M_2) by M_3 consists of

- (1) a biextension of \mathcal{B} of (G_1, G_2) by G_3 ;
- (2) a trivialization (= biaddictive section) Ψ_1 (resp. Ψ_2) of the biextension $(u_1, id_{G_2})^* \mathcal{B}$ (resp. $(id_{G_1}, u_2)^* \mathcal{B}$) of (X_1, G_2) by G_3 (resp. (G_1, X_2) by G_3) obtained as pull-back of the biextension \mathcal{B} via (u_1, id_{G_2}) (resp. (id_{G_1}, u_2));
- (3) a trivialization Ψ of the biextension $(u_1, u_2)^* \mathcal{B}$ of (X_1, X_2) by G_3 obtained as pull-back of the biextension \mathcal{B} by (u_1, u_2) , which coincides with the trivializations induced by Ψ_1 and Ψ_2 over $X_1 \times X_2$, i.e.

$$(u_1, id_{G_2})^* \Psi_2 = \Psi = (id_{G_1}, u_2)^* \Psi_1;$$

- (4) a morphism $\lambda : X_1 \times X_2 \rightarrow X_3$ of S -group schemes such that $u_3 \circ \lambda : X_1 \times X_2 \rightarrow G_3$ is compatible with the trivialization Ψ of the biextension $(u_1, u_2)^* \mathcal{B}$ of (X_1, X_2) by G_3 .

Let $M_i = [X_i \xrightarrow{u_i} G_i]$ and $M'_i = [X'_i \xrightarrow{u'_i} G'_i]$ (for $i = 1, 2, 3$) be 1-motives over S . Moreover let $(\mathcal{B}, \Psi_1, \Psi_2, \lambda)$ be a biextension of (M_1, M_2) by M_3 and let $(\mathcal{B}', \Psi'_1, \Psi'_2, \lambda')$ be a biextension of (M'_1, M'_2) by M'_3 .

Definition 1.1.2. A morphism of biextensions

$$(F, \Upsilon_1, \Upsilon_2, \Upsilon, g_3) : (\mathcal{B}, \Psi_1, \Psi_2, \lambda) \rightarrow (\mathcal{B}', \Psi'_1, \Psi'_2, \lambda')$$

consists of

- (1) a morphism $F = (F, f_1, f_2, f_3) : \mathcal{B} \rightarrow \mathcal{B}'$ from the biextension \mathcal{B} to the biextension \mathcal{B}' . In particular,

$$f_1 : G_1 \rightarrow G'_1 \quad f_2 : G_2 \rightarrow G'_2 \quad f_3 : G_3 \rightarrow G'_3$$

are morphisms of groups S -schemes.

(2) a morphism of biextensions

$$\Upsilon_1 = (\Upsilon_1, g_1, f_2, f_3) : (u_1, id_{G_2})^* \mathcal{B} \longrightarrow (u'_1, id_{G'_2})^* \mathcal{B}'$$

compatible with the morphism $F = (F, f_1, f_2, f_3)$ and with the trivializations Ψ_1 and Ψ'_1 , and a morphism of biextensions

$$\Upsilon_2 = (\Upsilon_2, f_1, g_2, f_3) : (id_{G_1}, u_2)^* \mathcal{B} \longrightarrow (id_{G'_1}, u'_2)^* \mathcal{B}'$$

compatible with the morphism $F = (F, f_1, f_2, f_3)$ and with the trivializations Ψ_2 and Ψ'_2 . In particular

$$g_1 : X_1 \longrightarrow X'_1 \quad g_2 : X_2 \longrightarrow X'_2$$

are morphisms of groups S -schemes.

(3) a morphism of biextensions $\Upsilon = (\Upsilon, g_1, g_2, f_3) : (u_1, u_2)^* \mathcal{B} \longrightarrow (u'_1, u'_2)^* \mathcal{B}'$ compatible with the morphism $F = (F, f_1, f_2, f_3)$ and with the trivializations Ψ and Ψ' .

(4) a morphism $g_3 : X_3 \longrightarrow X'_3$ of S -group schemes compatible with u_3 and u'_3 (i.e. $u'_3 \circ g_3 = f_3 \circ u_3$) and such that

$$\lambda' \circ (g_1 \times g_2) = g_3 \circ \lambda$$

Remark 1.1.3. The pair (g_3, f_3) defines a morphism of $\mathcal{M}(S)$ from M_3 to M'_3 . The pairs (g_1, f_1) and (g_2, f_2) define morphisms of $\mathcal{M}(S)$ from M_1 and M'_1 and from M_2 to M'_2 respectively.

We denote by $\mathbf{Biext}(M_1, M_2; M_3)$ the category of biextensions of (M_1, M_2) by M_3 . Like in the category of biextensions of semi-abelian schemes, in the category $\mathbf{Biext}(M_1, M_2; M_3)$ we can make the sum of two objects. Let $\text{Biext}^0(M_1, M_2; M_3)$ be the group of automorphisms of any biextension of (M_1, M_2) by M_3 , and let $\text{Biext}^1(M_1, M_2; M_3)$ be the group of isomorphism classes of biextensions of (M_1, M_2) by M_3 .

1.2. A more useful definition. From now on we will work on the topos \mathbf{T}_{fppf} associated to the site of locally of finite presentation S -schemes, endowed with the fppf topology.

Proposition [3] (10.2.14) furnishes a more symmetric description of 1-motives: consider the 7-uplet $(X, Y^\vee, A, A^*, v, v^*, \psi)$ where

- X and Y^\vee are two S -group schemes which are locally for the étale topology constant group schemes defined by finitely generated free \mathbb{Z} -modules. We have to think at X and at Y^\vee as character groups of S -tori which should be written (according to our notation) $X^\vee(1)$ and $Y(1)$, where X^\vee and Y are their cocharacter groups;
- A and A^* are two abelian S -schemes dual to each other;
- $v : X \longrightarrow A$ and $v^* : Y^\vee \longrightarrow A^*$ are two morphisms of S -group schemes; and
- ψ is a trivialization of the pull-back $(v, v^*)^* \mathcal{P}_A$ via (v, v^*) of the Poincaré biextension \mathcal{P}_A of (A, A^*) .

To have the data $(X, Y^\vee, A, A^*, v, v^*, \psi)$ is equivalent to have the 1-motive $M = [X \xrightarrow{u} G]$: In fact, to have the semi-abelian S -scheme G is the same thing as to have the morphism $v^* : Y^\vee \longrightarrow A^*$ (cf. [6] Exposé VIII 3.7, G corresponds to the biextension $(id_A, v^*)^* \mathcal{P}_A$ of (A, Y^\vee) by \mathbb{G}_m) and to have the morphism $u : X \longrightarrow$

G is equivalent to have the morphism $v : X \longrightarrow A$ and the trivialization ψ of $(v, v^*)^*\mathcal{P}_A$ (the trivialization ψ furnishes the lift $u : X \longrightarrow G$ of the morphism $v : X \longrightarrow A$).

Let $s : X \times Y^\vee \longrightarrow Y^\vee \times X$ be the morphism which permutates the factors. The 7-uplet $(Y^\vee, X, A^*, A, v^*, v, \psi \circ s)$ defines the so called **Cartier dual** of M : this is a 1-motive that we denote by M^* .

Remark 1.2.1. The pull-back $(v, v^*)^*\mathcal{P}_A$ by (v, v^*) of the Poincaré biextension \mathcal{P}_A of (A, A^*) is a biextension of (X, Y^\vee) by \mathbb{G}_m . According [5] Exposé X Corollary 4.5, we can suppose that the character group Y^\vee is constant, i.e. $\mathbb{Z}^{\text{rk} Y^\vee}$ (if necessary we localize over S for the étale topology). Moreover since by [6] Exposé VII (2.4.2) the category **Biext** is additive in each variable, we have that

$$\mathbf{Biext}(X, Y^\vee; \mathbb{G}_m) \cong \mathbf{Biext}(X, \mathbb{Z}; Y(1)).$$

We denote by

$$((v, v^*)^*\mathcal{P}_A) \otimes Y$$

the biextension of (X, \mathbb{Z}) by $Y(1)$ corresponding to the biextension $(v, v^*)^*\mathcal{P}_A$ through this equivalence of categories. The trivialization ψ of $(v, v^*)^*\mathcal{P}_A$ defines a trivialization

$$\psi \otimes Y$$

of $((v, v^*)^*\mathcal{P}_A) \otimes Y$, and vice versa.

We can now give a more useful definition of a biextension of two 1-motives by a third one:

Proposition 1.2.2. *Let $M_i = (X_i, Y_i^\vee, A_i, A_i^*, v_i, v_i^*, \psi_i)$ (for $i = 1, 2, 3$) be a 1-motive. A biextension $B = (B, \Psi'_1, \Psi'_2, \Psi', \Lambda)$ of (M_1, M_2) by M_3 consists of*

- (1) *a biextension of B of (A_1, A_2) by $Y_3(1)$;*
- (2) *a trivialization Ψ'_1 (resp. Ψ'_2) of the biextension $(v_1, id_{A_2})^*B$ (resp. $(id_{A_1}, v_2)^*B$) of (X_1, A_2) by $Y_3(1)$ (resp. of (A_1, X_2) by $Y_3(1)$) obtained as pull-back of the biextension B via (v_1, id_{A_2}) (resp. via (id_{A_1}, v_2));*
- (3) *a trivialization Ψ' of the biextension $(v_1, v_2)^*B$ of (X_1, X_2) by $Y_3(1)$ obtained as pull-back of the biextension B via (v_1, v_2) , which coincides with the trivializations induced by Ψ'_1 and Ψ'_2 over $X_1 \times X_2$, i.e.*

$$(v_1, id_{A_2})^*\Psi'_2 = \Psi' = (id_{A_1}, v_2)^*\Psi'_1;$$

- (4) *a morphism $\Lambda : (v_1, v_2)^*B \longrightarrow ((v_3, v_3^*)^*\mathcal{P}_{A_3}) \otimes Y_3$ of trivial biextensions, with $\Lambda|_{Y_3(1)}$ equal to the identity, such that the following diagram is commutative*

$$(1.2.1) \quad \begin{array}{ccc} Y_3(1) & = & Y_3(1) \\ \downarrow & & \downarrow \\ (v_1, v_2)^*B & \longrightarrow & ((v_3, v_3^*)^*\mathcal{P}_{A_3}) \otimes Y_3 \\ \Psi' \uparrow \downarrow & & \downarrow \psi_3 \otimes Y_3 \\ X_1 \times X_2 & \longrightarrow & X_3 \times \mathbb{Z}. \end{array}$$

Proof. According to the main Theorem of [2], to have the biextension B of (A_1, A_2) by $Y_3(1)$ is equivalent to have the biextension $\mathcal{B} = \iota_{3*}(\pi_1, \pi_2)^*B$ of (G_1, G_2) by G_3 , where for $i = 1, 2, 3$, $\pi_i : G_i \longrightarrow A_i$ is the projection of G_i over A_i and $\iota_i : Y_i(1) \longrightarrow G_i$ is the inclusion of $Y_i(1)$ over G_i .

The trivializations $(\Psi', \Psi'_1, \Psi'_2)$ and (Ψ, Ψ_1, Ψ_2) determine each others. By [2], both biextensions $(v_1, v_2)^*B$ and $((v_3, v_3^*)^*\mathcal{P}_{A_3}) \otimes Y_3$ are trivial. Hence to have the morphism of S -group schemes $\lambda : X_1 \times X_2 \longrightarrow X_3$ is equivalent to have the morphism of trivial biextensions $\Lambda : (v_1, v_2)^*B \longrightarrow ((v_3, v_3^*)^*\mathcal{P}_{A_3}) \otimes Y_3$ with $\Lambda|_{Y_3(1)}$ equal to the identity. In particular, through this equivalence λ corresponds to $\Lambda|_{X_1 \times X_2}$ and to require that $u_3 \circ \lambda : X_1 \times X_2 \longrightarrow G_3$ is compatible with the trivialization Ψ of $(u_1, u_2)^*\mathcal{B}$ corresponds to require the commutativity of the diagram (1.2.1). \square

Remark 1.2.3. The data (1), (2), (3) and (4) of definition 1.1.1 are equivalent respectively to the data (1), (2), (3) and (4) of Proposition 1.2.2.

1.3. Examples. We conclude this chapter giving some examples of biextensions of 1-motives by 1-motives. We will use the more useful definition of biextensions furnishes by Proposition 1.2.2.

(1) Let $M = [0 \longrightarrow A]$ be an abelian S -scheme with Cartier dual $M^* = [0 \longrightarrow A^*]$ and let $W(1)$ be an S -torus. A biextension of (M, M^*) by $W(1)$ is

$$(B, 0, 0, 0, 0)$$

where B is a biextension of (A, A^*) by $W(1)$. In particular **the Poincaré biextension of (M, M^*) by $\mathbb{Z}(1)$** is the biextension

$$(\mathcal{P}_A, 0, 0, 0, 0)$$

where \mathcal{P}_A is the Poincaré biextension of (A, A^*) .

(2) Let $M = (A, A^*, X, Y^\vee, v, v^*, \psi) = [X \xrightarrow{u} G]$ be a 1-motive over S and $M^* = [Y^\vee \xrightarrow{u^*} G^*]$ its Cartier dual. If \mathcal{P}_A denotes the Poincaré biextension of (A, A^*) , the semi-abelian S -scheme G (resp. G^*) corresponds to the biextension $(id_A, v^*)^*\mathcal{P}_A$ of (A, Y^\vee) by $\mathbb{Z}(1)$ (resp. $(v, id_{A^*})^*\mathcal{P}_A$ of (X, A^*) by \mathbb{G}_m) (cf. [6] Exposé VIII 3.7). **The Poincaré biextension of (M, M^*) by $\mathbb{Z}(1)$** is the biextension

$$(\mathcal{P}_A, \psi_1, \psi_2, \psi, 0)$$

where ψ_1 is the trivialization of the biextension $(id_A, v^*)^*\mathcal{P}_A$ which defines the morphism $u : X \longrightarrow G$, and ψ_2 is the trivialization of the biextension $(v, id_{A^*})^*\mathcal{P}_A$ which defines the morphism $u^* : Y^\vee \longrightarrow G^*$.

(3) Let $M = [X \xrightarrow{u} Y(1)]$ be the 1-motive without abelian part defined over S . Its Cartier dual is the 1-motive $M^* = [Y^\vee \xrightarrow{u^*} X^\vee(1)]$, where Y^\vee is the character group of the S -torus $Y(1)$ and $X^\vee(1)$ is the S -torus whose character group is X . **The Poincaré biextension of (M, M^*) by $\mathbb{Z}(1)$** is the biextension

$$(\underline{0}, 0, 0, \psi, 0)$$

where $\underline{0}$ is the trivial biextension of $(0, 0)$ by $\mathbb{Z}(1)$ and $\psi : X \times Y^\vee \longrightarrow \mathbb{Z}(1)$ is the biaddictive morphism defining $u : X \longrightarrow Y(1)$ and $u^* : Y^\vee \longrightarrow X^\vee(1)$. (We can view the trivialization ψ as a biaddictive morphism from $X \times Y^\vee$ to $\mathbb{Z}(1)$ because of the triviality of the biextension $\underline{0}$).

(4) Let $M = [X \xrightarrow{u} Y(1)]$ and $[V \xrightarrow{t} W(1)]$ be two 1-motives without abelian part defined over S , and let $M^* = [Y^\vee \xrightarrow{u^*} X^\vee(1)]$ be the Cartier dual of M . A biextension of (M, M^*) by $[V \xrightarrow{t} W(1)]$ is

$$(\underline{0}, 0, 0, \Psi, \lambda)$$

where $\underline{0}$ is the trivial biextension of $(0, 0)$ by $W(1)$, $\Psi : X \times Y^\vee \longrightarrow W(1)$ is a biaddictive morphism and $\lambda : X \times Y^\vee \longrightarrow V$ a morphism of S -group schemes, such that the diagram

$$(1.3.1) \quad \begin{array}{ccc} W(1) & = & W(1) \\ \Psi \uparrow & & \uparrow t \\ X \times Y^\vee & \xrightarrow{\lambda} & V \end{array}$$

is commutative. (We can view the trivialization Ψ as a biaddictive morphism from $X \times Y^\vee$ to $W(1)$ because of the triviality of the biextension $\underline{0}$).

(5) Let $M = [X \xrightarrow{u} Y(1)]$ be a 1-motive without abelian part defined over S , and $M^* = [Y^\vee \xrightarrow{u^*} X^\vee(1)]$ its Cartier dual. Let $[V \xrightarrow{t} A]$ be a 1-motive defined over S without toric part. A biextension of (M, M^*) by $[V \xrightarrow{t} A]$ is

$$(\underline{0}, 0, 0, 0, \lambda)$$

where $\underline{0}$ is the trivial biextension of $(0, 0)$ by 0 and $\lambda : X \times Y^\vee \longrightarrow V$ is a morphism of S -group schemes. Therefore

$$(1.3.2) \quad \mathbf{Biext}^1(M, M^*; [V \xrightarrow{t} A]) = \mathrm{Hom}_{S\text{-schemes}}(X \times Y^\vee, V).$$

(6) Let $M = [X \xrightarrow{u} A \times Y(1)] = (X, Y^\vee, A, A^*, v, v^* = 0, \psi)$ be a 1-motive defined over S , and $M^* = [Y^\vee \xrightarrow{u^*} G^*]$ its Cartier dual. Let $[V \xrightarrow{t} W(1)]$ be a 1-motive defined over S without abelian part. A biextension of (M, M^*) by $[V \xrightarrow{t} W(1)]$ is

$$(B, \Psi_1, \Psi_2, \Psi, \lambda)$$

where

- B is a biextension of (A, A^*) by $W(1)$;
- Ψ_1 (resp. Ψ_2) is a trivialization of the biextension $(v, id_{A^*})^*B$ (resp. $(id_A, v^*)^*B$) of (X, A^*) by $W(1)$ (resp. of (A, Y^\vee) by $W(1)$) obtained as pull-back of the biextension B via (v, id_{A^*}) (resp. via (id_A, v^*)). Since $v^* = 0$, the biextension $(id_A, v^*)^*B$ of (A, Y^\vee) by $W(1)$ is trivial;
- Ψ is a trivialization of the biextension $(v, v^*)^*B$ of (X, Y^\vee) by $W(1)$ obtained as pull-back of the biextension B via (v, v^*) , which coincides with the trivializations induced by Ψ_1 and Ψ_2 over $X \times Y^\vee$. Also the biextension $(v, v^*)^*B$ of (X, Y^\vee) by $W(1)$ is trivial and hence we can view Ψ as a biaddictive morphism from $X \times Y^\vee$ to $W(1)$;
- $\lambda : X \times Y^\vee \longrightarrow V$ is a morphism of S -group schemes such that the following diagram is commutative

$$(1.3.3) \quad \begin{array}{ccc} W(1) & = & W(1) \\ \Psi \uparrow & & \uparrow t \\ X \times Y^\vee & \longrightarrow & V. \end{array}$$

(7) Let $M = [X \xrightarrow{u} A \times Y(1)] = (X, Y^\vee, A, A^*, v, v^* = 0, \psi)$ be a 1-motive defined over S , and $M^* = [Y^\vee \xrightarrow{u^*} G^*]$ its Cartier dual. Let $[V \xrightarrow{t} A'] =$

$(V, 0, A', A'^*, t, 0, \psi')$ be a 1-motive defined over S without toric part. A biextension of (M, M^*) by $[V \xrightarrow{t} A']$ is

$$(\underline{\mathbf{0}}, 0, 0, \Psi, \lambda)$$

where $\underline{\mathbf{0}}$ is the trivial biextension of (A, A^*) by 0, $\Psi : X \times Y^\vee \longrightarrow X \times Y^\vee$ is a biaddictive morphism and $\lambda : X \times Y^\vee \longrightarrow V$ a morphism of S -group schemes, such that the diagram

$$(1.3.4) \quad \begin{array}{ccc} X \times Y^\vee & \xrightarrow{\lambda} & V \times \mathbb{Z} \\ \Psi \uparrow & & \uparrow \psi' \otimes 0 \\ X \times Y^\vee & \xrightarrow{\lambda} & V \times \mathbb{Z} \end{array}$$

is commutative. (We can view the trivialization Ψ as a biaddictive morphism from $X \times Y^\vee$ to $X \times Y^\vee$ because of the triviality of the biextension $\underline{\mathbf{0}}$ and for the meaning of $\psi' \otimes 0$ see (1.2.1)).

2. SOME MORPHISMS OF 1-MOTIVES

In this chapter we suppose that 1-motives over S generate a Tannakian category over a field K .

Let $\mathcal{M}(S)$ be the Tannakian category generated by 1-motives over S . The unit object of $\mathcal{M}(S)$ is the 1-motive $\mathbb{Z}(0) = [\mathbb{Z} \longrightarrow 0]$. We denote by $M^\vee \cong \underline{\text{Hom}}(M, \mathbb{Z}(0))$ the dual of the 1-motive M and $ev_M : M \otimes M^\vee \longrightarrow \mathbb{Z}(0)$ and $\delta_M : \mathbb{Z}(0) \longrightarrow M^\vee \otimes M$ the morphisms of $\mathcal{M}(S)$ which characterize M^\vee (cf. [4] (2.1.2)). The Cartier dual of M is the 1-motive

$$(2.0.5) \quad M^* = M^\vee \otimes \mathbb{Z}(1)$$

where $\mathbb{Z}(1)$ is the S -torus with cocharacter group \mathbb{Z} .

2.1. Motivic remarks about morphisms of 1-motives.

Lemma 2.1.1. *Let X_i (for $i = 1, 2, 3$) be a S -group scheme which is locally for the étale topology a constant group scheme defined by a finitely generated free \mathbb{Z} -module, and let G_i (for $i = 1, 2, 3$) be a semi-abelian S -scheme. In the category $\mathcal{M}(S)$, there are no morphisms from the tensor product $X_1 \otimes X_2$ to G , i.e.,*

$$\text{Hom}_{\mathcal{M}(S)}(X_1 \otimes X_2, G_3) = 0$$

and there are no morphisms from the tensor product $G_1 \otimes G_2$ to X_3 , i.e.,

$$\text{Hom}_{\mathcal{M}(S)}(G_1 \otimes G_2, X_3) = 0.$$

Proof. This lemma follows from the fact that morphisms of motives have to respect weights. In fact the pure motive $X_1 \otimes X_2$ has weight 0 and the mixed motive G has weight smaller or equal to -1, and the mixed motive $G_1 \otimes G_2$ has weight smaller or equal to -2 and the pure motive X_3 has weight 0. \square

Proposition 2.1.2. *Let G_i (for $i = 1, 2, 3$) be a semi-abelian S -scheme, i.e. an extension of an abelian S -scheme A_i by a S -torus $Y_i(1)$. We have that:*

$$(2.1.1) \quad \text{Hom}_{\mathcal{M}(S)}(G_1 \otimes G_2, Y_3(1)) = \text{Hom}_{\mathcal{M}(S)}(A_1 \otimes A_2, Y_3(1))$$

$$(2.1.2) \quad \text{Hom}_{\mathcal{M}(S)}(G_1 \otimes G_2, G_3) = \text{Hom}_{\mathcal{M}(S)}(G_1 \otimes G_2, Y_3(1))$$

Proof. The proof of these equalities is based on the fact that morphisms of motives have to respect weights.

For $i = 1, 2, 3$, denote by $\pi_i : G_i \rightarrow A_i$ the projection of G_i over A_i and $\iota_i : Y_i(1) \rightarrow G_i$ the inclusion of $Y_i(1)$ in G_i . Thanks to the projection (π_1, π_2) , each morphism $A_1 \otimes A_2 \rightarrow Y_3(1)$ can be lifted to a morphism from $G_1 \otimes G_2$ to $Y_3(1)$. In the other hand, since the motive $Y_1(1) \otimes Y_2(1)$ has weight -4 , each morphism $G_1 \otimes G_2 \rightarrow Y_3(1)$ factorizes through $A_1 \otimes A_2$.

Through the inclusion ι_3 , each morphism $G_1 \otimes G_2 \rightarrow Y_3(1)$ defines a morphism $G_1 \otimes G_2 \rightarrow G_3$. In the other hand, since the motive $G_1 \otimes G_2$ has weight less or equal to -2 , each morphism $G_1 \otimes G_2 \rightarrow G_3$ factorizes through $Y_3(1)$. \square

Therefore we have:

Corollary 2.1.3. *In $\mathcal{M}(S)$, a morphism from the tensor product of two semi-abelian S -schemes to a semi-abelian S -scheme is a morphism from the tensor product of the underlying abelian S -schemes to the underlying S -torus:*

$$\mathrm{Hom}_{\mathcal{M}(S)}(G_1 \otimes G_2, G_3) = \mathrm{Hom}_{\mathcal{M}(S)}(A_1 \otimes A_2, Y_3(1)).$$

2.2. Morphisms from a finite tensor products of 1-motives to a 1-motive.

Definition 2.2.1. In the category $\mathcal{M}(S)$, the morphism $M_1 \otimes M_2 \rightarrow M_3$ from the tensor product of two 1-motives to a third 1-motive is an isomorphism class of biextensions of (M_1, M_2) by M_3 (cf. (1.1.1)). We define

$$(2.2.1) \quad \mathrm{Hom}_{\mathcal{M}(S)}(M_1 \otimes M_2, M_3) = \mathrm{Biext}^1(M_1, M_2; M_3)$$

In other words, the biextensions of two 1-motives by a 1-motive are the “geometrical interpretation” of the morphisms of $\mathcal{M}(S)$ from the tensor product of two 1-motives to a 1-motive.

Remark 2.2.2. The set $\mathrm{Hom}_{\mathcal{M}(S)}(M_1 \otimes M_2, M_3)$ is a group.

Remark 2.2.3. The confrontation of the main Theorem of [4] with Corollary 2.1.3, shows that this definition is compatible with what we know about morphisms between 1-motives.

Definition 2.2.4. Let M_i and M'_i (for $i = 1, 2, 3$) be 1-motives over S . The notion 1.1.2 of morphisms of biextensions defines a **morphism from the group of morphisms of $\mathcal{M}(S)$ from $M_1 \otimes M_2$ to M_3 , to the group of morphisms of $\mathcal{M}(S)$ from $M'_1 \otimes M'_2$ to M'_3** , i.e.

$$\mathrm{Hom}_{\mathcal{M}(S)}(M_1 \otimes M_2, M_3) \rightarrow \mathrm{Hom}_{\mathcal{M}(S)}(M'_1 \otimes M'_2, M'_3).$$

Remark 2.2.5. Using the notations of 1.1.2, if we denote b the morphism $M_1 \otimes M_2 \rightarrow M_3$ corresponding to the biextension $(\mathcal{B}, \Psi_1, \Psi_2, \lambda)$ of (M_1, M_2) by M_3 and by b' the morphism $M'_1 \otimes M'_2 \rightarrow M'_3$ corresponding to the biextension $(\mathcal{B}', \Psi'_1, \Psi'_2, \lambda')$ of (M'_1, M'_2) by M'_3 , the morphism

$$(F, \Upsilon_1, \Upsilon_2, \Upsilon, g_3) : (\mathcal{B}, \Psi_1, \Psi_2, \lambda) \rightarrow (\mathcal{B}', \Psi'_1, \Psi'_2, \lambda')$$

of biextensions defines the vertical arrows of the following diagram of morphisms of $\mathcal{M}(S)$

$$\begin{array}{ccc}
M_1 \otimes M_2 & \xrightarrow{b} & M_3 \\
\downarrow & & \downarrow \\
M'_1 \otimes M'_2 & \xrightarrow{b'} & M'_3.
\end{array}$$

It is clear now why from the data $(F, \Upsilon_1, \Upsilon_2, \Upsilon, g_3)$ we get a morphism of $\mathcal{M}(S)$ from M_3 to M'_3 as remarked in 1.1.3. Moreover since $M_1 \otimes \mathbb{Z}(0)$, $M'_1 \otimes \mathbb{Z}(0)$, $\mathbb{Z}(0) \otimes M_2$ and $\mathbb{Z}(0) \otimes M'_2$, are sub-1-motives of the motives $M_1 \otimes M_2$ and $M'_1 \otimes M'_2$, it is clear that from the data $(F, \Upsilon_1, \Upsilon_2, \Upsilon, g_3)$ we get morphisms from M_1 to M'_1 and from M_2 to M'_2 as remarked in 1.1.3.

Lemma 2.2.6. *Let l and i be positive integers and let $M_j = [X_j \xrightarrow{u_j} G_j]$ (for $j = 1, \dots, l$) be a 1-motive defined over S . Denote by M_0 or X_0 the 1-motive $\mathbb{Z}(0) = [\mathbb{Z} \rightarrow 0]$. If $i \geq 1$ and $l + 1 \geq i$, the motive $\otimes_{j=1}^l M_j / W_{-i}(\otimes_{j=1}^l M_j)$ is isogeneous to the motive*

$$(2.2.2) \quad \sum \left(\otimes_{k \in \{\nu_0, \dots, \nu_{l-i+1}\}} X_k \right) \bigotimes \left(\otimes_{j \in \{\iota_0, \dots, \iota_{i-1}\}} M_j / W_{-i}(\otimes_{j \in \{\iota_0, \dots, \iota_{i-1}\}} M_j) \right)$$

where the sum is taken over all the $(l-i+1)$ -uplets $\{\nu_0, \dots, \nu_{l-i+1}\}$ and all the $(i-1)$ -uplets $\{\iota_0, \dots, \iota_{i-1}\}$ of $\{0, 1, \dots, l\}$ such that $\{\nu_0, \dots, \nu_{l-i+1}\} \cap \{\iota_0, \dots, \iota_{i-1}\} = \emptyset$, $\nu_0 < \nu_1 < \dots < \nu_{l-i+1}$, $\iota_0 < \iota_1 < \dots < \iota_{i-1}$, $\nu_a \neq \nu_b$ and $\iota_c \neq \iota_d$, for all $a, b \in \{0, \dots, l-i+1\}$, $a \neq b$ and $c, d \in \{0, \dots, i-1\}$, $c \neq d$.

Proof. 1-motives M_j are composed by pure motives of weight 0 (the lattice part X_j), -1 (the abelian part A_j) and -2 (the toric part $Y_j(1)$). Consider the pure motive $\text{Gr}_{-i}^W(\otimes_{j=1}^l M_j)$: it is a finite sum of tensor products of l factors of weight 0, -1 other -2. If $i = l$ the tensor product

$$A_1 \otimes A_2 \otimes \dots \otimes A_l$$

contains no factors of weight 0. For each i strictly bigger than l , it is also easy to construct a tensor product of l factors whose total weight is $-i$ and in which no factor has weight 0 (for example if $i = l + 2$ we take

$$Y_1(1) \otimes Y_2(1) \otimes A_3 \otimes \dots \otimes A_l.)$$

However if i is strictly smaller than l , in each of these tensor products of l factors, there is at least one factor of weight 0, i.e. one of the X_j for $j = 1, \dots, l$.

Now fix a i strictly smaller than l . The tensor products where there are less factors of weight 0 are exactly those where there are more factors of weight -1. Hence in the pure motive $\text{Gr}_{-i}^W(\otimes_{j=1}^l M_j)$, the tensor products with less factors of weight 0 are of the type

$$X_{\nu_1} \otimes \dots \otimes X_{\nu_{l-i}} \otimes A_{\iota_1} \otimes \dots \otimes A_{\iota_i}$$

Thanks to these observations, the conclusion is clear.

Remark that we have only an isogeny because in the 1-motive (2.2.2) the factor

$$X_{\nu_1} \otimes X_{\nu_2} \otimes \dots \otimes X_{\nu_p} \otimes \mathcal{Y}_{\iota_1} \otimes \mathcal{Y}_{\iota_2} \otimes \dots \otimes \mathcal{Y}_{\iota_{l-p}}$$

appears with multiplicity “ $p+m$ ” where m is the number of \mathcal{Y}_{ι_q} (for $q = 1, \dots, l-p$) which are of weight 0, instead of appearing only once like in the 1-motive $\otimes_j M_j / W_{-i}(\otimes_j M_j)$. In particular for each i we have that

$$\begin{aligned} \mathrm{Gr}_0^W \left(\sum (\otimes_k X_k) \otimes (\otimes_j M_j / W_{-i}) \right) &= l \, \mathrm{Gr}_0^W \left(\otimes_j M_j / W_{-i} \right) \\ \mathrm{Gr}_{-1}^W \left(\sum (\otimes_k X_k) \otimes (\otimes_j M_j / W_{-i}) \right) &= (l-1) \, \mathrm{Gr}_{-1}^W \left(\otimes_j M_j / W_{-i} \right) \end{aligned}$$

□

We will denote by $\mathcal{M}^{\mathrm{iso}}(S)$ the Tannakian category generated by the iso-1-motives, i.e. by 1-motives modulo isogenies.

Theorem 2.2.7. *Let M and M_1, \dots, M_l be 1-motives over S . In the category $\mathcal{M}^{\mathrm{iso}}(S)$, the morphism $\otimes_{j=1}^l M_j \rightarrow M$ from a finite tensor product of 1-motives to a 1-motive is the sum of copies of isomorphism classes of biextensions of (M_i, M_j) by M for $i, j = 1, \dots, l$ and $i \neq j$. We have that*

$$\mathrm{Hom}_{\mathcal{M}^{\mathrm{iso}}(S)}(\otimes_{j=1}^l M_j, M) = \sum_{\substack{i, j \in \{1, \dots, l\} \\ i \neq j}} \mathrm{Biext}^1(M_i, M_j; M)$$

Proof. Because morphisms between motives have to respect weights, the non trivial components of the morphism $\otimes_{j=1}^l M_j \rightarrow M$ are the one of the morphism

$$\otimes_{j=1}^l M_j / W_{-3}(\otimes_{j=1}^l M_j) \rightarrow M.$$

Using the equality obtained in Lemma 2.2.6 with $i = -3$, we can write explicitly this last morphism in the following way

$$\sum_{\substack{\iota_1 < \iota_2 \text{ and } \nu_1 < \dots < \nu_{l-2} \\ \iota_1, \iota_2 \notin \{\nu_1, \dots, \nu_{l-2}\}}} X_{\nu_1} \otimes \dots \otimes X_{\nu_{l-2}} \otimes (M_{\iota_1} \otimes M_{\iota_2} / W_{-3}(M_{\iota_1} \otimes M_{\iota_2})) \rightarrow M.$$

Since “to tensorize a motive by a motive of weight 0” means to take a certain number of copies of the motive, from definition 2.2.1 we get the expected conclusion. □

2.3. Examples. Now we give some examples of morphisms from the tensor product of two 1-motives to a 1-motive. According to our definition 2.2.1 of morphisms of $\mathcal{M}(S)$, these examples are in parallel with the examples 1.3.

(1) Let $M = [0 \rightarrow A]$ be an abelian S -scheme with Cartier dual $M^* = [0 \rightarrow A^*]$ and let $W(1)$ be an S -torus. By definition, the biextension $(B, 0, 0, 0, 0)$ of (M, M^*) by $W(1)$ is a morphism of $\mathcal{M}(S)$ from $A \otimes A^*$ to $W(1)$. In particular, the Poincaré biextension \mathcal{P}_A of (A, A^*) by $\mathbb{Z}(1)$ is the classical **Weil pairing of A**

$$\mathcal{P}_A : A \otimes A^* \rightarrow \mathbb{Z}(1).$$

(2) Let M be a 1-motive over S and M^* its Cartier dual. The Poincaré biextension \mathcal{P}_M of (M, M^*) by $\mathbb{Z}(1)$ (cf. 1.3) is the so called **Weil pairing of M**

$$\mathcal{P}_M : M \otimes M^* \rightarrow \mathbb{Z}(1),$$

which expresses the Cartier duality between M and M^* .

The valuation map $ev_M : M \otimes M^\vee \rightarrow \mathbb{Z}(0)$ of M expresses the duality between M and M^\vee as objects of the Tannakian category $\mathcal{M}(S)$. By (2.0.5) the evaluation map ev_M is the twist by $\mathbb{Z}(-1)$ of the Weil pairing of M :

$$ev_M = \mathcal{P}_M \otimes \mathbb{Z}(-1) : M \otimes M^\vee \rightarrow \mathbb{Z}(0).$$

(3) Let M be the 1-motive without abelian part $[X \xrightarrow{u} Y(1)]$ defined over S . Its Cartier dual is the 1-motive $M^* = [Y^\vee \xrightarrow{u^*} X^\vee(1)]$. We will compute the

morphism from the tensor product $M \otimes M^*$ to the torus $\mathbb{Z}(1)$, which is by definition the isomorphism class of the Poincaré biextension of (M, M^*) by $\mathbb{Z}(1)$. Regarding M and M^* as complexes of commutative groups S -schemes concentrated in degree 0 and -1, their tensor product is the complex

$$(2.3.1) \quad [X \otimes Y^\vee \xrightarrow{(-id_X \otimes u^*, u \otimes id_{Y^\vee})} X \otimes X^\vee(1) + Y(1) \otimes Y^\vee \xrightarrow{u \otimes id_{X^\vee} + id_{Y(1)} \otimes u^*} Y(1) \otimes X^\vee(1)]$$

where $X \otimes Y^\vee$ is a pure motive of weight 0, $X \otimes X^\vee(1) + Y(1) \otimes Y^\vee$ is a pure motive of weight -2 and $Y(1) \otimes X^\vee(1)$ is a pure motive of weight -4. Because of the weights, the only non-trivial component of a morphism from the tensor product $M \otimes M^*$ to the torus $\mathbb{Z}(1)$ is

$$(2.3.2) \quad X \otimes X^\vee(1) + Y(1) \otimes Y^\vee \longrightarrow \mathbb{Z}(1)$$

By example 1.3 (3), the Poincaré biextension of (M, M^*) by $\mathbb{Z}(1)$ is the biextension $(0, 0, 0, \psi, 0)$, where $\psi : X \times Y^\vee \longrightarrow \mathbb{Z}(1)$ is the biaddictive morphism defining $u : X \longrightarrow Y(1)$ and $u^* : Y^\vee \longrightarrow X^\vee(1)$. This biaddictive morphism ψ defines the only non-trivial component (2.3.2) of $M \otimes M^* \longrightarrow \mathbb{Z}(1)$ through the following commutative diagram

$$\begin{array}{ccc} X \otimes Y^\vee & & \\ \downarrow (-id_X \otimes u^*, u \otimes id_{Y^\vee}) & \searrow \psi & \\ X \otimes X^\vee(1) + Y(1) \otimes Y^\vee & \longrightarrow & \mathbb{Z}(1) \end{array}$$

(4) Let $M = [X \xrightarrow{u} Y(1)]$ and $[V \xrightarrow{t} W(1)]$ be two 1-motives without abelian part defined over S , and let $M^* = [Y^\vee \xrightarrow{u^*} X^\vee(1)]$ be the Cartier dual of M . According (2.3.1), the only non trivial components of a morphism from the tensor product $M \otimes M^*$ to $[V \xrightarrow{t} W(1)]$ are

$$(2.3.3) \quad X \otimes Y^\vee \longrightarrow V$$

$$(2.3.4) \quad X \otimes X^\vee(1) + Y(1) \otimes Y^\vee \longrightarrow W(1)$$

By example 1.3 (4), a biextension of (M, M^*) by $[V \xrightarrow{t} W(1)]$ is $(\underline{0}, 0, 0, \Psi, \lambda)$ where $\underline{0}$ is the trivial biextension of $(0, 0)$ by $W(1)$, $\Psi : X \times Y^\vee \longrightarrow W(1)$ is a biaddictive morphism and $\lambda : X \times Y^\vee \longrightarrow V$ a morphism of S -group schemes, such that the diagram (1.3.1) is commutative. The morphism λ defines the non-trivial component (2.3.3) between motives of weight 0. Through the commutative diagram

$$\begin{array}{ccc} X \otimes Y^\vee & & \\ \downarrow (-id_X \otimes u^*, u \otimes id_{Y^\vee}) & \searrow \Psi & \\ X \otimes X^\vee(1) + Y(1) \otimes Y^\vee & \longrightarrow & W(1) \end{array}$$

the morphism Ψ defines the non-trivial component (2.3.4) between motives of weight -2. The commutativity of the diagram (1.3.1) impose the compatibility between these two non trivial components.

(5) Let $M = [X \xrightarrow{u} Y(1)]$ be a 1-motive without abelian part defined over S , and $M^* = [Y^\vee \xrightarrow{u^*} X^\vee(1)]$ its Cartier dual. Let $[V \xrightarrow{t} A]$ a 1-motive defined over

S without toric part. Because of the weights, according (2.3.1) the only non trivial component of a morphism from the tensor product $M \otimes M^*$ to $[V \xrightarrow{t} A]$ is

$$(2.3.5) \quad X \otimes Y^\vee \longrightarrow V$$

By example 1.3 (4), a biextension of (M, M^*) by $[V \xrightarrow{t} A]$ is $(\underline{0}, 0, 0, \lambda)$ where $\underline{0}$ is the trivial biextension of $(0, 0)$ by 0 and $\lambda : X \times Y^\vee \longrightarrow V$ is a morphism of S -group schemes. This morphism λ defines the only non trivial component (2.3.5) of the morphism $M \otimes M^* \longrightarrow [V \xrightarrow{t} A]$. In particular by (1.3.2)

$$\mathrm{Hom}_{\mathcal{M}(S)}(M \otimes M^*, [V \xrightarrow{t} A]) = \mathrm{Hom}_{S\text{-schemes}}(X \times Y^\vee, V).$$

(6) Let $M = [X \xrightarrow{u} A \times Y(1)] = (X, Y^\vee, A, A^*, v, v^* = 0, \psi)$ be a 1-motive defined over S , and $M^* = [Y^\vee \xrightarrow{u^*} G^*]$ its Cartier dual. Let $[V \xrightarrow{t} W(1)]$ be a 1-motive defined over S without abelian part. Because of the weights, the only non trivial components of a morphism from the tensor product $M \otimes M^*$ to $[V \xrightarrow{t} W(1)]$ are

$$(2.3.6) \quad X \otimes Y^\vee \longrightarrow V$$

$$(2.3.7) \quad A \otimes A^* + X \otimes X^\vee(1) + Y(1) \otimes Y^\vee \longrightarrow W(1)$$

By example 1.3 (6), a biextension of (M, M^*) by $[V \xrightarrow{t} W(1)]$ is $(B, \Psi_1, \Psi_2, \Psi, \lambda)$ where B is a biextension of (A, A^*) by $W(1)$, $\Psi : X \times Y^\vee \longrightarrow W(1)$ is a biadditive morphism and $\lambda : X \times Y^\vee \longrightarrow V$ is a morphism of S -group schemes such that the diagram (1.3.3) is commutative. The morphism λ defines the non-trivial component (2.3.6) between motives of weight 0. Through the commutative diagram

$$\begin{array}{ccc} X \otimes Y^\vee & & \\ \downarrow & \searrow \Psi & \\ A \otimes A^* + X \otimes X^\vee(1) + Y(1) \otimes Y^\vee & \longrightarrow & W(1) \end{array}$$

the morphism Ψ and the biextension B define the non-trivial component (2.3.7) between motives of weight -2: to be more precise, the biextension B is the component $A \otimes A^* \longrightarrow W(1)$ and the morphism Ψ determines the component $X \otimes X^\vee(1) + Y(1) \otimes Y^\vee \longrightarrow W(1)$. The commutativity of the diagram (1.3.3) imposes the compatibility between these two non trivial components. Remark that the factor $A \otimes A^*$ plays no role in the diagram (1.3.3) because $v^* = 0$.

(7) Let $M = [X \xrightarrow{u} A \times Y(1)] = (X, Y^\vee, A, A^*, v, v^* = 0, \psi)$ be a 1-motive defined over S , and $M^* = [Y^\vee \xrightarrow{u^*} G^*]$ its Cartier dual. Let $[V \xrightarrow{t} A'] = (V, 0, A', A'^*, t, 0, \psi')$ be a 1-motive defined over S without toric part. Because of the weights, the only non trivial components of a morphism from the tensor product $M \otimes M^*$ to $[V \xrightarrow{t} A']$ are

$$(2.3.8) \quad X \otimes Y^\vee \longrightarrow V$$

$$(2.3.9) \quad X \otimes A^* + A \otimes Y^\vee(1) \longrightarrow A'$$

By example 1.3 (7), a biextension of (M, M^*) by $[V \xrightarrow{t} A]$ is $(\underline{0}, 0, 0, \Psi, \lambda)$ where $\underline{0}$ is the trivial biextension of (A, A^*) by 0, $\Psi : X \times Y^\vee \longrightarrow X \times Y^\vee$ is a biadditive morphism and $\lambda : X \times Y^\vee \longrightarrow V$ a morphism of S -group schemes, such that the

diagram (1.3.4) is commutative. The morphism λ defines the component (2.3.8) between motives of weight 0 and the biaddictive morphism Ψ determines the component (2.3.9). The commutativity of the diagram (1.3.4) imposes the compatibility between these two non trivial components.

REFERENCES

- [1] C. Bertolin, *Le radical unipotent du groupe de Galois motivique d'un 1-motif*, Math. Ann. 327 (2003).
- [2] C. Bertolin, *Biextensions of semi-abelian schemes by semi-abelian schemes*, in preparation (2003).
- [3] P. Deligne, *Théorie de Hodge III*, Pub. Math. de l'I.H.E.S 44 (1975).
- [4] P. Deligne, *Catégories tannakiennes*, The Grothendieck Festschrift II, Birkhäuser 87 (1990).
- [5] SGA3 II: Schémas en groupes, LN 152 (1970).
- [6] SGA7 I: Groupes de Monodromie en Géométrie Algébrique, LN 288 (1972).

NWF I -MATHEMATIK, UNIVERSITÄT REGENSBURG, 93040 REGENSBURG, GERMANY
E-mail address: `cristiana.bertolin@mathematik.uni-regensburg.de`